

Some identities of Frobenius-Euler polynomials arising from Frobenius-Euler basis

by

Dae San Kim and Taekyun Kim

Abstract

In this paper, we give some new and interesting identities which are derived from the basis of Frobenius-Euler. Recently, Simsek et als(see [13]) have given some identities of q -analogue of Frobenius-Euler polynomials related to q -Bernstein polynomials. From the methods of our paper, we can also derive the results and identities of Simsek et als (cf.[13]).

1 Introduction

Let $\lambda(\neq 1) \in \mathbf{C}$. As is well known, the Frobenius-Euler polynomials are defined by the generating function to be

$$\frac{1-\lambda}{e^t-\lambda}e^{xt} = e^{H(x|\lambda)t} = \sum_{n=0}^{\infty} H_n(x|\lambda) \frac{t^n}{n!}, \quad (1)$$

with the usual convention about replacing $H^n(x|\lambda)$ by $H_n(x|\lambda)$ (see [1–6]). In the special case, $x=0$, $H_n(0|\lambda) = H_n(\lambda)$ are called the n -th Frobenius-Euler numbers.

Thus, by(1), we get

$$(H(\lambda) + 1)^n - \lambda H_n(\lambda) = H_n(1|\lambda) - \lambda H_n(\lambda) = (1-\lambda)\delta_{0,n}, \quad (2)$$

where $\delta_{0,n}$ is the Kronecker symbol.

From (1), we can derive the following equation :

$$H_n(x|\lambda) = (H(\lambda) + x)^n = \sum_{l=0}^n \binom{n}{l} H_{n-l}(\lambda) x^l, \quad (\text{see [6–13]}) . \quad (3)$$

Thus, by (3), we easily see that the leading coefficient of $H_n(x|\lambda)$ is $H_0(\lambda) = 1$. So, $H_n(x|\lambda)$ is a monic polynomials of degree n with coefficients in $\mathbf{Q}(\lambda)$. From (1), we have

$$\sum_{n=0}^{\infty} (H_n(x+1|\lambda) - \lambda H_n(x|\lambda)) \frac{t^n}{n!} = \frac{(1-\lambda)e^{(x+1)t}}{e^t - \lambda} - \lambda \frac{1-\lambda}{e^t - \lambda} e^{xt}. \quad (4)$$

Thus, by (4), we get

$$H_n(x+1|\lambda) - \lambda H_n(x|\lambda) = (1-\lambda)x^n, \text{ for } n \in \mathbf{Z}_+. \quad (5)$$

It is easy to show that

$$\frac{d}{dx} H_n(x|\lambda) = \frac{d}{dx} (H(\lambda) + x)^n = n H_{n-1}(x|\lambda), \quad (n \in \mathbf{N}). \quad (6)$$

From (6), we have

$$\int_0^1 H_n(x|\lambda) dx = \frac{1}{n+1} (H_{n+1}(1|\lambda) - H_{n+1}(\lambda)) = \frac{\lambda-1}{n+1} H_{n+1}(\lambda). \quad (7)$$

Let $\mathbb{P}_n(\lambda) = \{p(x) \in \mathbf{Q}(\lambda)[x] \mid \deg p(x) \leq n\}$ be a vector space over $\mathbf{Q}(\lambda)$. Then we note that $\{H_0(x|\lambda), H_1(x|\lambda), \dots, H_n(x|\lambda)\}$ is a good basis for $\mathbb{P}_n(\lambda)$. In this paper, we develop some new methods to obtain some new identities and properties of Frobenius-Euler polynomials which are derived from the basis of the Frobenius-Euler polynomials. Those methods are useful in studying the identities of Frobenius-Euler polynomials.

2 Some identities of Frobenius-Euler polynomials

Let us take $p(x) \in \mathbb{P}_n(\lambda)$. Then $p(x)$ can be expressed as a $\mathbf{Q}(\lambda)$ -linear combination of $H_0(x|\lambda), \dots, H_n(x|\lambda)$ as follows :

$$p(x) = b_0 H_0(x|\lambda) + b_1 H_1(x|\lambda) + \dots + b_n H_n(x|\lambda) = \sum_{k=0}^n b_k H_k(x|\lambda). \quad (8)$$

Let us define the operator Δ_λ by

$$g(x) = \Delta_\lambda p(x) = p(x+1) - \lambda p(x). \quad (9)$$

From (9), we can derive the following equation (10) :

$$g(x) = \triangle_\lambda p(x) = \sum_{k=0}^n b_k (H_k(x+1|\lambda) - \lambda H_k(x|\lambda)) = (1-\lambda) \sum_{k=0}^n b_k x^k. \quad (10)$$

For $r \in \mathbf{Z}_+$, let us take the r -th derivative of $g(x)$ in (10) as follows :

$$g^{(r)}(x) = (1-\lambda) \sum_{k=r}^n k(k-1) \cdots (k-r+1) b_k x^{k-r}, \text{ where } g^{(r)}(x) = \frac{d^r g(x)}{dx^r}. \quad (11)$$

Thus, by (11), we get

$$g^{(r)}(0) = \frac{d^r g(x)}{dx^r} \Big|_{x=0} = (1-\lambda) r! b_r. \quad (12)$$

From (12), we have

$$b_r = \frac{g^{(r)}(0)}{(1-\lambda) r!} = \frac{1}{(1-\lambda) r!} (p^{(r)}(1) - \lambda p^{(r)}(0)), \quad (13)$$

where $r \in \mathbf{Z}_+$, and $p^{(r)}(0) = \frac{d^r p(x)}{dx^r} \Big|_{x=0}$. Therefore, by (13), we obtain the following theorem.

Theorem 1. For $\lambda (\neq 1) \in \mathbf{C}$, $n \in \mathbf{Z}_+$,
let $p(x) \in \mathbb{P}_n(\lambda)$ with $p(x) = \sum_{k=0}^n b_k H_k(x|\lambda)$. Then we have

$$b_k = \frac{1}{(1-\lambda) k!} g^{(k)}(0) = \frac{1}{(1-\lambda) k!} (p^{(k)}(1) - \lambda p^{(k)}(0)).$$

Let us take $p(x) = H_n(x|\lambda^{-1})$. Then, by Theorem 1, we get

$$H_n(x|\lambda^{-1}) = \sum_{k=0}^n b_k H_k(x|\lambda), \quad (14)$$

where

$$\begin{aligned} b_k &= \frac{1}{(1-\lambda) k!} \frac{n!}{(n-k)!} \{H_{n-k}(1|\lambda^{-1}) - \lambda H_{n-k}(\lambda^{-1})\} \\ &= \frac{1}{1-\lambda} \binom{n}{k} \{H_{n-k}(1|\lambda^{-1}) - \lambda H_{n-k}(\lambda^{-1})\} \\ &= \frac{1}{1-\lambda} \binom{n}{k} \{(1-\lambda^{-1}) 0^{n-k} + \frac{1}{\lambda} H_{n-k}(\lambda^{-1}) - \lambda H_{n-k}(\lambda^{-1})\}. \end{aligned} \quad (15)$$

By (14) and (15), we get

$$\begin{aligned}
& H_n(x|\lambda^{-1}) \\
&= -\frac{1}{\lambda}H_n(x|\lambda) + \sum_{k=0}^n \left\{ \frac{\binom{n}{k}}{\lambda(1-\lambda)} H_{n-k}(\lambda^{-1}) - \frac{\lambda \binom{n}{k}}{1-\lambda} H_{n-k}(\lambda^{-1}) H_k \right\}(x|\lambda) \\
&= -\frac{1}{\lambda}H_n(x|\lambda) + \sum_{k=0}^n \binom{n}{k} \frac{1+\lambda}{\lambda} H_{n-k}(\lambda^{-1}) H_k(x|\lambda).
\end{aligned} \tag{16}$$

Therefore, by (16), we obtain the following theorem.

Theorem 2. *For $n \in \mathbf{Z}_+$, we have*

$$\lambda H_n(x|\lambda^{-1}) + H_n(x|\lambda) = (1+\lambda) \sum_{k=0}^n \binom{n}{k} H_{n-k}(\lambda^{-1}) H_k(x|\lambda).$$

Let

$$p(x) = \sum_{k=0}^n H_k(x|\lambda) H_{n-k}(x|\lambda) \in \mathbb{P}_n(\lambda). \tag{17}$$

From Theorem 2, we note that $p(x)$ can be generated by $\{H_0(x|\lambda), H_1(x|\lambda), \dots, H_n(x|\lambda)\}$ as follows:

$$p(x) = \sum_{k=0}^n H_k(x|\lambda) H_{n-k}(x|\lambda) = \sum_{k=0}^n b_k H_k(x|\lambda). \tag{18}$$

By (17), we get

$$p^{(k)}(x) = \frac{(n+1)!}{(n-k+1)!} \sum_{l=k}^n H_{l-k}(x|\lambda) H_{n-k}(x|\lambda), \tag{19}$$

and

$$\begin{aligned}
b_k &= \frac{1}{(1-\lambda)k!} \{p^{(k)}(1) - \lambda p^{(k)}(0)\} \\
&= \frac{(n+1)!}{(1-\lambda)k!(n-k+1)!} \sum_{l=k}^n \{H_{l-k}(1|\lambda)H_{n-l}(1|\lambda) - \lambda H_{l-k}(\lambda)H_{n-l}(\lambda)\} \\
&= \frac{n+1}{(1-\lambda)(n-k+1)} \binom{n}{k} \sum_{l=k}^n \{(\lambda H_{l-k}(\lambda) + (1-\lambda)\delta_{0,l-k})(\lambda H_{n-l} + \\
&\quad (1-\lambda)\delta_{0,n-l}) - \lambda H_{l-k}(\lambda)H_{n-l}(\lambda)\} \\
&= \frac{n+1}{(1-\lambda)(n-k+1)} \binom{n}{k} \sum_{l=k}^n \{\lambda(1-\lambda)\delta_{0,l-k}H_{n-l}(\lambda) + \lambda(1-\lambda) \\
&\quad \times H_{l-k}(\lambda)\delta_{0,n-l} + (1-\lambda)^2\delta_{0,l-k}\delta_{0,n-l} + \lambda(\lambda-1)H_{l-k}(\lambda)H_{n-l}(\lambda)\} \\
&= \frac{n+1}{(1-\lambda)(n-k+1)} \binom{n}{k} \sum_{l=k}^n \{\lambda(\lambda-1)H_{l-k}(\lambda)H_{n-l}(\lambda) + \lambda(1-\lambda) \\
&\quad \times H_{n-k}(\lambda) + \lambda(1-\lambda)H_{n-k}(\lambda) + (1-\lambda)^2\delta_{n,k}\} \\
&= \frac{n+1}{n-k+1} \binom{n}{k} \sum_{l=k}^n \{-\lambda H_{l-k}(\lambda)H_{n-l}(\lambda) + 2\lambda H_{n-k}(\lambda) + (1-\lambda)\delta_{n,k}\}.
\end{aligned} \tag{20}$$

From (18) and (20), we have

$$\begin{aligned}
\sum_{k=0}^n H_k(x|\lambda)H_{n-k}(x|\lambda) &= (n+1) \sum_{k=0}^{n-1} \frac{\binom{n}{k}}{n-k+1} \sum_{l=k}^n \{(-\lambda)H_{l-k}(\lambda)H_{n-l}(\lambda) \\
&\quad + 2\lambda H_{n-k}(\lambda)\} H_k(x|\lambda) + (n+1)H_n(x|\lambda).
\end{aligned} \tag{21}$$

Therefore, by (21), we obtain the following theorem

Theorem 3. For $n \in \mathbf{Z}_+$, we have

$$\begin{aligned}
&\frac{1}{n+1} \sum_{k=0}^n H_k(x|\lambda)H_{n-k}(x|\lambda) \\
&= \sum_{k=0}^{n-1} \frac{\binom{n}{k}}{n-k+1} \sum_{l=k}^n \{(-\lambda)H_{l-k}(\lambda)H_{n-l}(\lambda) + 2\lambda H_{n-k}(\lambda)\} H_k(x|\lambda) + H_n(x|\lambda).
\end{aligned}$$

Let us consider

$$p(x) = \sum_{k=0}^n \frac{1}{k!(n-k)!} H_k(x|\lambda) H_{n-k}(x|\lambda) \in \mathbb{P}_n(\lambda). \quad (22)$$

By Theorem 1, $p(x)$ can be expressed by

$$p(x) = \sum_{k=0}^n b_k H_k(x|\lambda). \quad (23)$$

From (22), we have

$$p^{(r)}(x) = 2^r \sum_{k=r}^n \frac{H_{k-r}(x|\lambda) H_{n-k}(x|\lambda)}{(k-r)!(n-k)!}, (r \in \mathbf{Z}_+). \quad (24)$$

By Theorem 1, we get

$$\begin{aligned} b_k &= \frac{1}{2k!} \{p^{(k)}(1) - p^{(k)}(0)\} \\ &= \frac{2^{k-1}}{k!} \sum_{l=k}^n \frac{1}{(l-k)!(n-l)!} \{H_{l-k}(1|\lambda) H_{n-l}(1|\lambda) - \lambda H_{l-k}(\lambda) H_{n-l}(\lambda)\} \\ &= \frac{2^{k-1}}{k!} \sum_{l=k}^n \frac{1}{(l-k)!(n-l)!} \{(\lambda H_{l-k}(\lambda) + (1-\lambda)\delta_{0,l-k})(\lambda H_{n-l}(\lambda) \\ &\quad + (1-\lambda)\delta_{0,n-l}) - \lambda H_{l-k}(\lambda) H_{n-l}(\lambda)\} \\ &= \frac{2^{k-1}}{k!} \left\{ \sum_{l=k}^n \frac{\lambda(\lambda-1) H_{l-k}(\lambda) H_{n-l}(\lambda)}{(l-k)!(n-l)!} + \frac{2\lambda(1-\lambda) H_{n-k}(\lambda)}{(n-k)!} + (1-\lambda)^2 \delta_{n,k} \right\} \\ &= \begin{cases} \frac{2^{k-1}}{k!} \sum_{l=k}^n \left\{ \frac{\lambda(\lambda-1) H_{l-k}(\lambda) H_{n-l}(\lambda)}{(l-k)!(n-l)!} + \frac{2\lambda(1-\lambda) H_{n-k}(\lambda)}{(n-k)!} \right\}, & \text{if } k \neq n \\ \frac{2^{n-1}(1-\lambda)}{n!}, & \text{if } k = n \end{cases} \end{aligned} \quad (25)$$

Therefore, by (25), we obtain the following theorem.

Theorem 4. For $n \in \mathbf{Z}_+$, we have

$$\begin{aligned} &\sum_{k=0}^n \frac{1}{k!(n-k)!} H_k(x|\lambda) H_{n-k}(x|\lambda) \\ &= \sum_{k=0}^{n-1} \frac{2^{k-1}}{k!} \sum_{l=k}^n \left\{ \frac{\lambda(\lambda-1) H_{l-k}(\lambda) H_{n-l}(\lambda)}{(l-k)!(n-l)!} + \frac{2\lambda(1-\lambda) H_{n-k}(\lambda)}{(n-k)!} \right\} H_k(x|\lambda) \\ &\quad + \frac{2^{n-1}(1-\lambda)}{n!} H_n(x|\lambda). \end{aligned}$$

3 Higher-order Frobenius-Euler polynomials

For $n \in \mathbf{Z}_+$, the Frobenius-Euler polynomials of order r are defined by the generating function to be

$$\begin{aligned} \left(\frac{1-\lambda}{e^t-\lambda}\right)^r e^{xt} &= e^{H^{(r)}(x|\lambda)t} \\ &= \sum_{n=0}^{\infty} H_n^{(r)}(x|\lambda) \frac{t^n}{n!}, \end{aligned} \quad (26)$$

with the usual convention about replacing $(H^{(r)}(x|\lambda))^n$ by $H_n^{(r)}(x|\lambda)$, (see [1–10]). In the special case, $x=0$, $H_n^{(r)}(0|\lambda) = H_n^{(r)}(\lambda)$ are called the n -th Frobenius-Euler numbers of order r , (see [8–9]).

From (26), we have

$$H_n^{(r)}(x|\lambda) = (H^{(r)}(\lambda) + x)^n = \sum_{l=0}^n \binom{n}{l} H_{n-l}^{(r)}(\lambda) x^l, \quad (27)$$

with the usual convention about replacing $(H^{(r)}(\lambda))^n$ by $H_n^{(r)}(\lambda)$.

By (26), we get

$$H_n^{(r)}(\lambda) = \sum_{n_1+\dots+n_r=n} \binom{n}{n_1, n_2, \dots, n_r} H_{n_1}(\lambda) \cdots H_{n_r}(\lambda), \quad (28)$$

where $\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!}$. From (27) and (28), we note that the leading coefficient of $H_n^{(r)}(x|\lambda)$ is given by

$$\begin{aligned} H_0^{(r)}(\lambda) &= \sum_{n_1+\dots+n_r=0} \binom{n}{n_1, n_2, \dots, n_r} H_{n_1}(\lambda) \cdots H_{n_r}(\lambda) \\ &= H_0(\lambda) \cdots H_0(\lambda) = 1. \end{aligned} \quad (29)$$

Thus, by (29), we see that $H_n^{(r)}$ is a monic polynomial of degree n with coefficients in $\mathbf{Q}(\lambda)$. From (26), we have

$$H_n^{(0)}(x|\lambda) = x^n, \text{ for } n \in \mathbf{Z}_+, \quad (30)$$

and

$$\frac{\partial}{\partial x} H_n^{(r)}(x|\lambda) = \frac{\partial}{\partial x} (H^{(r)}(\lambda) + x)^n = n H_{n-1}^{(r)}(x|\lambda), \quad (r \geq 0). \quad (31)$$

It is not difficult to show that

$$H_n^{(r)}(x+1|\lambda) - \lambda H_n^{(r)}(x|\lambda) = (1-\lambda)H_n^{(r-1)}(x|\lambda). \quad (32)$$

Now, we note that $\{H_0^{(r)}(x|\lambda), H_1^{(r)}(x|\lambda), \dots, H_n^{(r)}(x|\lambda)\}$ is also a good basis for $\mathbb{P}_n(\lambda)$.

Let us define the operator D as $Df(x) = \frac{df(x)}{dx}$ and let $p(x) \in \mathbb{P}_n(\lambda)$. Then $p(x)$ can be written as

$$p(x) = \sum_{k=0}^n C_k H_k^{(r)}(x|\lambda). \quad (33)$$

From (9) and (32), we have

$$\Delta_\lambda H_n^{(r)}(x|\lambda) = H_n^{(r)}(x+1|\lambda) - \lambda H_n^{(r)}(x|\lambda) = (1-\lambda)H_n^{(r-1)}(x|\lambda). \quad (34)$$

Thus, by (33) and (34), we get

$$\Delta_\lambda^r p(x) = (1-\lambda)^r \sum_{k=0}^n C_k H_k^{(0)}(x|\lambda) = (1-\lambda)^r \sum_{k=0}^n C_k x^k. \quad (35)$$

Let us take the k -th derivative of $\Delta_\lambda^r p(x)$ in (35).

Then we have

$$D^k(\Delta_\lambda^r p(x)) = (1-\lambda)^r \sum_{l=k}^n \frac{l!}{(l-k)!} C_l x^{l-k}. \quad (36)$$

Thus, from (36), we have

$$D^k(\Delta_\lambda^r p(0)) = (1-\lambda)^r \sum_{l=k}^n \frac{l! C_l}{(l-k)!} 0^{l-k} = (1-\lambda)^r k! C_k. \quad (37)$$

Thus, by (37), we get

$$\begin{aligned} C_k &= \frac{D^k(\Delta_\lambda^r p(0))}{(1-\lambda)^r k!} \\ &= \frac{\Delta_\lambda^r(D^k p(0))}{(1-\lambda)^r k!} = \frac{1}{(1-\lambda)^r k!} \sum_{j=0}^r \binom{r}{j} (-\lambda)^{(r-j)} D^k p(j). \end{aligned} \quad (38)$$

Therefore, by (33) and (38), we obtain the following theorem.

Theorem 5. For $r \in \mathbf{Z}_+$, let $p(x) \in \mathbb{P}_n(\lambda)$ with

$$p(x) = \frac{1}{(1-\lambda)^r} \sum_{k=0}^n C_k H_k^{(r)}(x|\lambda), \quad (C_k \in \mathbf{Q}(\lambda)).$$

Then we have

$$C_k = \frac{1}{(1-\lambda)^r k!} \sum_{j=0}^r \binom{r}{j} (-\lambda)^{r-j} D^k p(j).$$

That is,

$$p(x) = \frac{1}{(1-\lambda)^r} \sum_{k=0}^n \left(\sum_{j=0}^r \frac{1}{k!} \binom{r}{j} (-\lambda)^{r-j} D^k p(j) \right) H_k^{(r)}(x|\lambda).$$

Let us take $p(x) = H_n(x|\lambda) \in \mathbf{P}_n(\lambda)$. Then, by Theorem 5, $p(x) = H_n(x|\lambda)$ can be generated by $\{H_0^{(r)}(x|\lambda), H_1^{(r)}(x|\lambda), \dots, H_n^{(r)}(x|\lambda)\}$ as follows :

$$H_n(x|\lambda) = \sum_{k=0}^n C_k H_k^{(r)}(x|\lambda), \quad (39)$$

where

$$C_k = \frac{1}{(1-\lambda)^r} \frac{1}{k!} \sum_{j=0}^r \binom{r}{j} (-\lambda)^{r-j} D^k p(j), \quad (40)$$

and

$$p^{(k)}(x) = D^k p(x) = n(n-1) \cdots (n-k+1) H_{n-k}(x|\lambda) = \frac{n!}{(n-k)!} H_{n-k}(x|\lambda). \quad (41)$$

By (40) and (41), we get

$$C_k = \frac{1}{(1-\lambda)^r} \binom{n}{k} \sum_{j=0}^r \binom{r}{j} (-\lambda)^{r-j} H_{n-k}(j|\lambda). \quad (42)$$

Therefore, by (39) and (42), we obtain the following theorem.

Theorem 6. For $n \in \mathbf{Z}_+$, we have

$$H_n(x|\lambda) = \frac{1}{(1-\lambda)^r} \sum_{k=0}^n \binom{n}{k} \left(\sum_{j=0}^r \binom{r}{j} (-\lambda)^{r-j} H_{n-k}(j|\lambda) \right) H_k^{(r)}(x|\lambda).$$

Let us assume that $p(x) = H_n^{(r)}(x|\lambda)$.
Then we have

$$\begin{aligned} p^k(x) &= n(n-1)\cdots(n-k+1)H_{n-k}^{(r)}(x|\lambda) \\ &= \frac{n!}{(n-k)!}H_{n-k}^{(r)}(x|\lambda). \end{aligned} \quad (43)$$

From Theorem 1, we note that $p(x) = H_n^{(r)}(x|\lambda)$ can be expressed as a linear combination of $H_0(x|\lambda), H_1(x|\lambda), \dots, H_n(x|\lambda)$:

$$H_n^{(r)}(x|\lambda) = \sum_{k=0}^n b_k H_k(x|\lambda), \quad (44)$$

where

$$\begin{aligned} b_k &= \frac{1}{(1-\lambda)k!} \{p^k(1) - \lambda p^{(k)}(0)\} \\ &= \frac{n!}{(1-\lambda)k!(n-k)!} \{H_{n-k}^{(r)}(1|\lambda) - \lambda H_{n-k}^{(r)}(\lambda)\}. \end{aligned} \quad (45)$$

By (34) and (45), we get

$$b_k = \frac{1}{(1-\lambda)^2} \binom{n}{k} H_{n-k}^{(r-1)}(\lambda). \quad (46)$$

Therefore, by (44) and (46), we obtain the following theorem.

Theorem 7. *For $n \in \mathbf{Z}_+$, we have*

$$H_n^{(r)}(x|\lambda) = \frac{1}{(1-\lambda)^2} \sum_{k=0}^n \binom{n}{k} H_{n-k}^{(r-1)}(\lambda) H_k(x|\lambda).$$

References

- [1] M. Can, M. Cenkci, V. Kurt, Y. Simsek, *Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Euler L-functions*, Adv. Stud. Contemp. Math., 18(2009), no. 2, 135-160.
- [2] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w - q -Geonocchi numbers* Adv. Stud. Contemp. Math. 19(2009), no. 1, 39-57.

- [3] L. Carlitz, *A note on q -Eulerian numbers*, J. Combin. Theory Ser. A 25(1978), no. 1, 90–94.
- [4] L. C. Jang, *A study on the distribution of twisted q -Genocchi polynomials*, Adv. Stud. Contemp. Math. 18(2009), no. 2, 181-189.
- [5] D. S. Kim, D.V. Dolgy, T. Kim, S.-H. Rim, *Some formulae for the product of two Bernoulli and Euler polynomials*, Abstr. Appl. Anal. 2012(2012), Article ID 784307, 15 pages.
- [6] T. Kim, J. Choi, *A note on the product of Frobenius-Euler polynomials arising from the p -adic integral on $\mathbb{Z}_p$* , Adv. Stud. Contemp. Math. 22 (2012), no. 2, 215-223.
- [7] T. Kim, *An identity of the symmetry for the Frobenius-Euler polynomials associated with the fermionic p -adic invariant q -integrals on $\mathbb{Z}_p$* , Rocky Mountain J. Math. 41 (2011), no. 1, 239-247.
- [8] T. Kim, *Identities involving Frobenius-Euler polynomials arising from non-linear differential equations*, J. Number Theory 132(2012), no. 12, 2854–2865.
- [9] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p -adic invariant integral on $\mathbb{Z}_p$* , Russ. J. Math. Phys. 16(2009), no. 1, 93-96.
- [10] S.-H. Rim, J. Jeong, *On the modified q -Euler numbers of higher order with weight*, Adv. Stud. Contemp. Math. 22(2012), no. 1, 93-98.
- [11] S.-H. Rim, J. Lee, *Some identities on the twisted (h, q) -Genocchi numbers and polynomials associated with q -Bernstein polynomials*, Int. J. Math. Math. Sci. 2011(2011), Art. ID 482840, 8 pages.
- [12] C. S. Ryoo, *A note on the Frobenius-Euler polynomials*, Proc. Jangjeon Math. Soc. 14(2011), no. 4, 495–501.
- [13] Y. Simsek, A. Bayad, V. Loksha, *q -Bernstein polynomials related to q -Frobenius-Euler polynomials, l -functions, and q -Stirling numbers*, Math. Methods Appl. Sci. 35(2012), no. 8, 877-884.
- [14] Y. Simsek, O. Yurekli, V. Kurt, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math., 15(2007), no. 2, 187-194.

- [15] K. Shiratani, *On the Euler numbers*, Mem. Fac. Sci., Kyushu Univ., Ser. A 27(1973), 1-5.
- [16] K. Shiratani, S. Yamamoto, *On a p -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci. Kyushu Univ. Ser. A 39(1985), no. 1, 113-125.

Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

e-mail: dskim@sogang.ac.kr

Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

e-mail: tkkim@kw.ac.kr